

ATTACHMENT 1 - Heat Exhaustion and Heat Stroke

[Better Health Channel: Heat stress and heat related illness](#)

Heat Illness

Heat exhaustion starts slowly and if not quickly treated can progress to heatstroke. In heatstroke, a child's temperature reaches 105° F (40.5° C) or higher. Heatstroke requires **immediate emergency medical care** and can be fatal.



Signs and Symptoms

Of heat exhaustion:

- increased thirst
- weakness
- fainting
- muscle cramps
- nausea and vomiting
- irritability
- headache
- increased sweating
- cool, clammy skin
- elevation of body temperature to less than 105° F (40.5° C)

Of heatstroke:

- severe headache
- weakness, dizziness
- confusion
- rapid breathing and heartbeat
- loss of consciousness leading to coma
- seizures
- may not be sweating
- flushed, hot, dry skin
- elevation of body temperature to 105° F (40.5° C) or higher



What to Do:

For a child with symptoms of heatstroke, **seek emergency medical care immediately**. In cases of heat exhaustion or while awaiting help for a child with possible heatstroke:

- Bring the child indoors or into the shade immediately.
- Undress the child.
- Have the child lie down; elevate feet slightly.
- If the child is alert, place in cool bath water. If outside, spray the child with mist from a garden hose.
- If the child is alert and coherent, give frequent sips of cool, clear fluids.
- If the child is vomiting, turn onto his or her side to prevent choking.



Think Prevention!

- Teach kids to always drink plenty of fluids before and during any activity in hot, sunny weather – even if they aren't thirsty.
- Make sure kids wear light-colored, loose clothing.
- Do not have your child participate in heavy activity outdoors during the hottest hours of the day.
- Teach kids to come indoors immediately whenever they feel overheated.